

Computation in Coxeter groups II. Minimal roots

Bill Casselman
 Mathematics Department
 University of British Columbia
 Canada
 cass@math.ubc.ca

Abstract. *In the recent paper (Casselman, 2001) I described how a number of ideas due to Fokko du Cloux and myself could be incorporated into a reasonably efficient program to carry out multiplication in arbitrary Coxeter groups. At the end of that paper I discussed how this algorithm could be used to build the reflection table of minimal roots, which could in turn form the basis of a much more efficient multiplication algorithm. In this paper, following a suggestion of Bob Howlett, I explain how results due to Brigitte Brink can be used to construct the minimal root reflection table directly and more efficiently.*

Introduction

A **Coxeter system** is a pair (W, S) where W is a group with a set of generators S and relations

$$(st)^{m_{s,t}} = 1$$

for pairs s, t in S . These groups play an important role in mathematics far from visible in this simple definition. Much work has been done on them, particularly on finite and affine Weyl groups, but many phenomena involving Coxeter groups remain unexplained, and it is likely that computer explorations will be even more significant in the future than they have been so far. These groups become extremely complex as the size of S grows, however, and computer programs dealing with them must be extremely efficient to be useful. Conventional symbolic algebra packages normally fail to handle the difficulties satisfactorily. This paper is the second of a series in which I describe what may be even in the long term the most efficient algorithms to do basic computations in arbitrary Coxeter groups, at least on serial machines. These algorithms depend strongly on mathematical results of Brigitte Brink, Robert Howlett, and Fokko du Cloux.

The first problem one encounters is how to multiply two elements of the group, which is by no means a simple matter. For familiar examples such as S_n there are many satisfactory solutions, and for the Weyl groups of Kac-Moody algebras one can use the representation of the group on the root lattice to reduce many problems to integer arithmetic. But arbitrary groups, even sometimes the extraordinary finite groups H_3 and H_4 , are computationally more demanding. Even with Weyl groups integer overflow may occur when working with the root lattice. In addition, solving computational problems in this domain leads to mathematically interesting questions as well.

The best combinatorial solutions—certainly, those of greatest theoretical interest—seem to be those in which elements of the group are represented in terms of products of elements of S . More explicitly, impose an ordering on S . Suppose w to be equal to the product $s_1 s_2 \dots s_n$ as a product of generators in S , and for each i let w_i be the partial product $s_1 \dots s_i$. This expression for w is called its **InverseShortLex normal form** if for each i the element s_i is least in S such that $\ell(w_i s_i) < \ell(w_i)$. It is the **shortest** representation of w least in **lexicographical order** when read **in reverse**. The multiplication problem can now be posed: *given the normal form of w , what is that of sw ?* The first general solution was implicit in work published by Jacques Tits in the mid 1960's. It had the virtue of being entirely combinatorial in nature, working only with the normal form of w , but it was extremely

inefficient, requiring both time and storage space roughly exponentially proportional, or worse, to the length n of the normal form of w .

A second solution was proposed by Fokko du Cloux around 1990, but explicitly described only for finite Coxeter groups. It was this that I worked out in more detail in (Casselman, 2001). It is mildly recursive in nature, of unknown theoretical complexity but in practice not unreasonably slow. It also has the tremendous virtue of requiring little machine memory. If one wants to be able to carry out just a few multiplications, it is likely that this method cannot be improved on. But if one wants to carry out a great many multiplications for a given Coxeter group, then it is probably best to build an auxiliary data structure that will make multiplication much simpler. A structure perfectly suited to this purpose is a table describing how the generators of S act on the **minimal roots** of (Brink and Howlett, 1993). As will be recalled later on, it can be used in a multiplication algorithm of complexity essentially linear in the length of w , allowing multiplication to take place without backtracking in a left-to-right scan of the normal form of w .

In the paper (Casselman, 2001) I proposed that du Cloux's algorithm could be used as a kind of bootstrap to build this reflection table, but as I mentioned at the end of that paper it is not difficult to find groups where the number of minimal roots is rather small, hence the table itself also fairly small, but the task of building it in this way extremely tedious. This is unsatisfactory.

After I explained this method to Robert Howlett, he and I experimented with other tools, and after a short while he suggested that the results of Brigitte Brink's thesis seemed to be applicable. In this thesis, almost all of which has been summarized in her published articles, she describes rather explicitly what the set of minimal roots looks like for any Coxeter group. However, it was not at all apparent at first glance, at least to me, that what she does is computationally practical. Howlett showed that my first impression was false by producing a program in the symbolic algebra package *Magma* that builds the minimal root reflection table of an arbitrary Coxeter group. This implementation was much slower than subsequent programs written by Fokko du Cloux and myself, but already dealt easily with a previously difficult group described in my paper. It might be worthwhile to say that the new algorithm depends on several crucial results of Brink's thesis, but not on her more or less explicit lists, interesting though they may be.

The principal drawback of the new algorithm as the basis of multiplication is that it claims an amount of machine memory proportional to the number of minimal roots. The number of minimal roots can grow quite fast with the size of S , so this is not a negligible objection. It thus exemplifies the usual programming trade-off between time and space. Familiar Coxeter groups are deceptive in this regard—for classical series of finite and affine groups the number of minimal roots is roughly $|S|^2$, whereas one of the first really difficult groups Howlett's program dealt with was an exotic Coxeter group with $|S| = 22$ and several hundred thousand minimal roots. (Howlett's *Magma* program took 69 minutes to handle it, he tells me, whereas a program in *C* written by Fokko du Cloux took 10 seconds on a comparable machine.) Of course, such groups are so complicated that there is not much exploration one can expect to do with them anyway, and in practice the number of minimal roots is only a relatively minor impediment.

In the rest of this paper I will first recall the basic properties of minimal roots; explain how they are to be used in multiplication; summarize results of Brigitte Brink's thesis in the manner most useful for my purposes; and finally explain how to apply these results in practice to construct the minimal root reflection table. I should say that what I am going to come up with is somewhat different from Howlett's program, and also quite different from a related program of Fokko du Cloux. It uses a few more advanced features of Brink's

thesis than Howlett's program. It is less sophisticated than du Cloux's program, but has the advantage of being possibly clearer in conception.

There is not much originality in this paper. Once Howlett had suggested using the results of Brink's thesis, the path to a practical program was probably almost determined. On the other hand, this path did not appear obvious to me until after a great deal of experiment, and I feel it will be useful to place here a record of the outcome. I should also say that the order of exposition and its occasional geometric emphasis are new. I will be satisfied if this exposition awakens interest in the work of Brink, Howlett, and du Cloux. This whole area of research is one of great charm, but it has apparently and unfortunately awakened little interest at large. This is likely due to the fact that it lies somewhere in the great no man's land between pure mathematics and practical computation—*terra incognita* on most maps, even in the 21st century. I must also record that both my programs and this exposition have benefited enormously from conversations with Fokko du Cloux. As I learned long ago to my own chagrin, competing directly in programming with him can be an embarrassing experience, but working alongside him is, on the contrary, at once pleasant and educational.

1. Minimal roots

The algorithms to be described are intimately related to geometric realizations of Coxeter groups. Following Brink and Howlett, I shall work here exclusively with the **standard realization**. To each element s of S is associated an element α_s of a basis Δ of a real vector space V . An inner product is defined on V according to the formula

$$\alpha_s \bullet \alpha_t = -\cos(\pi/m_{s,t}) = -(\zeta_{2m} + \zeta_{2m}^{-1})/2 \quad (m = m_{s,t}, \zeta_{2m} = e^{2\pi i/2m}).$$

In particular, $\alpha_s \bullet \alpha_s = 1$, $\alpha_s \bullet \alpha_t = -1$ whenever s and t generate an infinite group, and $-1 < \alpha_s \bullet \alpha_t \leq 1/2$ whenever they generate a finite non-abelian group. A representation of W on V is determined by mapping elements of S to reflections:

$$sv = v - 2(v \bullet \alpha_s)\alpha_s.$$

Recall that the **Coxeter graph** has as nodes the elements of S , and an edge linking s and t if $m_{s,t}$, which is called the **degree** of the edge, is more than 2. In other words, the edges link s and t with $\alpha_s \bullet \alpha_t \neq 0$. In case $\alpha \bullet \beta \neq 0$, I write $\alpha \sim \beta$. A **simple link** in the graph is one with degree 3, a **multiple link** one of higher degree. The formula for reflection amounts to the specification that if $\lambda = \sum \lambda_\gamma \gamma$ then under reflection by s_α all the coordinates of λ remain constant except that indexed by α , which changes to

$$(s\lambda)_\alpha = -\lambda_\alpha + \sum_{\alpha \sim \beta} [-2\alpha \bullet \beta] \lambda_\beta.$$

In calculations, it is frequently best to work with $2\alpha \bullet \beta$ rather than the dot-product itself, since it is at once simpler and more efficient to do so.

This representation is faithful. Let

$$C = \{v \mid v \bullet \alpha_s > 0\}.$$

The faces of this simplicial cone are the closures of the open simplices

$$C_T = \{v \in V \mid \langle \alpha, v \rangle = 0 \text{ for } \alpha \in T, \langle \alpha, v \rangle > 0 \text{ for } \alpha \notin T\}$$

where $T \subseteq S$. Thus C itself is $C_\emptyset$, and a face of codimension one is some C_α . The union of the domains $w\bar{C}$ is a convex conical region called the **Tits cone** $\mathcal{C}$ of the realization. The

group W acts discretely on $\mathcal{C}$, and the closure $\overline{\mathcal{C}}$ of the open simplicial cone is a fundamental domain for this action. Each face of $w\overline{\mathcal{C}}$ is the transform of a unique $\overline{\mathcal{C}}_T$; it is said to be **labeled** by T .

A transform $w\alpha_s$ of an element of Δ is called a **root** of this realization. The root hyperplane $\lambda = 0$ is fixed by the **root reflection** s_λ where $s_{w\alpha} = ws_\alpha w^{-1}$, the conjugate of an element in S . For any root λ , define

$$\begin{aligned}\mathcal{C}_{\lambda=0} &= \{v \in \mathcal{C} \mid \langle \lambda, v \rangle = 0\} \\ \mathcal{C}_{\lambda \geq 0} &= \{v \in \mathcal{C} \mid \langle \lambda, v \rangle \geq 0\} \\ \mathcal{C}_{\lambda \leq 0} &= \{v \in \mathcal{C} \mid \langle \lambda, v \rangle \leq 0\}.\end{aligned}$$

A root λ is called **positive** if $\lambda > 0$ on $\mathcal{C}$, **negative** if $\lambda < 0$ on $\mathcal{C}$. Every root is either positive or negative, which means that no root hyperplane ever intersects $\mathcal{C}$. A root is positive if and only if it can be expressed $\lambda = \sum_{\alpha \in \Delta} \lambda_\alpha \alpha$ with $\lambda_\alpha \geq 0$.

1.1. Proposition. *Suppose that λ and μ are positive roots. The following are equivalent:*


- (a) $\lambda \bullet \mu = \cos(\pi k / m_{s,t})$ with $0 < k < m_{s,t}$ for some s, t in S ;
- (b) $|\lambda \bullet \mu| < 1$;
- (c) the group generated by the root reflections s_λ and s_μ is finite;
- (d) the hyperplanes $\lambda = 0$ and $\mu = 0$ intersect in the interior of $\mathcal{C}$.

PROOF. This is implicit in (Vinberg, 1971). That (a) implies (b) is trivial; that (b) implies (c) is a consequence of the simplest part of his discussion of pairs of reflections. If G is any finite subgroup of W then the G -orbit of any point in the interior of $\mathcal{C}$ will be finite, and since $\mathcal{C}$ is convex the centre of mass of the orbit will be in the interior of $\mathcal{C}$ as well. This shows that (c) implies (d). Suppose $L = \{\lambda = 0\} \cap \{\mu = 0\}$ to contain a point in the interior of $\mathcal{C}$. The whole of $\mathcal{C}$ is tiled by simplices wC_T . Points in the open cones wC are fixed only by 1 in W , and those in some wC_α are fixed only by 1 and a single reflection. Therefore the linear space L must contain some wC_T with $|T| = 2$. This means that s and t can be simultaneously conjugated into W_T , and shows that (d) implies (a). $\square$

1.2. Corollary. *If λ and μ are distinct positive roots such that $|\lambda \bullet \mu| \geq 1$, then either $\mathcal{C}_{\lambda \geq 0}$ contains $\mathcal{C}_{\mu \geq 0}$ or $\mathcal{C}_{\mu \geq 0}$ contains $\mathcal{C}_{\lambda \geq 0}$.*

PROOF. If the intersection of $\lambda = 0$ and $\mu = 0$ with the interior of $\mathcal{C}$ is null, then the sign of μ on the intersection of $\lambda = 0$ with that interior will be constant. Suppose it is negative. I claim that the region $\mathcal{C}_{\mu \geq 0}$ is contained in the region $\mathcal{C}_{\lambda \geq 0}$. If not, suppose that P is a point of $\mathcal{C}$ with $\langle \mu, P \rangle \geq 0$ and $\langle \lambda, P \rangle < 0$. If P_0 is any point in the interior of $\mathcal{C}$, then the open line segment (P_0, P) is contained in the interior of $\mathcal{C}$, it is also contained in the region $\mathcal{C}_{\mu \geq 0}$, and it must intersect the hyperplane $\lambda = 0$. At that point μ will be non-negative, a contradiction. If the sign is positive, switch λ and μ . $\square$

Following Brink and Howlett, I say that the positive root λ **dominates** the positive root μ if the region $\mathcal{C}_{\lambda \geq 0}$ contains the region $\mathcal{C}_{\mu \geq 0}$. Another way of putting this is to require that $\lambda > 0$ in every chamber wC on which $\mu > 0$. Loosely speaking, a **minimal root** is one which dominates only itself. It is an important and remarkable fact about arbitrary Coxeter groups, first proved by Brink and Howlett, that the number of minimal roots is always finite. The significance of this result does not appear in familiar cases, since all roots are minimal when W is finite, and the minimal roots for affine Coxeter groups are rather simple in nature.



In the rest of this section I recall from (Brink and Howlett, 1993) some basic facts about minimal roots. Proofs and statements will often be the same as theirs, and I'll refer to them when this happens. But in some places what I take from them is only implicit in their paper, and in other places I have given both statements and proofs a geometrical flavour, somewhat as in (Casselman, 1995).

1.3. Lemma. (B & H 2.2(ii)) *Suppose λ and μ to be positive roots with λ dominating μ . If $w\mu$ is positive, then $w\lambda$ is also positive, and dominates $w\mu$.*

PROOF. Under the hypothesis, $\mathcal{C}_{w\lambda \geq 0}$ contains $\mathcal{C}_{w\mu \geq 0}$. If $w\mu$ is positive, the second contains C , and therefore $w\lambda > 0$ on C as well. $\square$

It is well known that s in S permutes the complement of α_s in the set of all positive roots, or in other words that if s is in S and λ a positive root, then either $\lambda = \alpha_s$ and $s\lambda < 0$, or $s\lambda > 0$. For the minimal roots, there is a similar range of options:

1.4. Proposition. *If λ is a minimal root and s in S then exactly one of these three options holds:*

- $\lambda = \alpha_s$ and $s\lambda < 0$;
- $s\lambda$ is again minimal;
- $s\lambda$ dominates α_s .

PROOF. If $\lambda \neq \alpha > 0$, then $s_\alpha \lambda > 0$. If it is not minimal, say it dominates $\beta > 0$ with $s_\alpha \lambda \neq \beta$. If $\beta \neq \alpha$, then $s_\alpha \beta > 0$, and by the previous Lemma λ dominates $s_\alpha \beta$. Since λ is minimal, this can only happen if $\lambda = s_\alpha \beta$, a contradiction. $\square$

The **minimal root reflection table** is a matrix ρ of size $|S| \times N$, where N is the number of minimal roots. The entries are either minimal roots, or **virtual** minimal roots I arbitrarily label as $\ominus$ and $\oplus$. I set $\rho(s, \lambda) = \mu$ if $s\lambda = \mu$ is minimal, $\rho(s, \lambda) = \ominus$ if $\lambda = \alpha_s$, and $\rho(s, \lambda) = \oplus$ if $s\lambda$ is a non-minimal positive root. This table is one of the fundamental data structures of a Coxeter group.

The **depth** $\delta(\lambda)$ of a positive root λ is the length of the smallest w such that $w^{-1}\lambda < 0$. Equivalently, it is one less than the length of the shortest gallery $C = C_0, C_1, \dots, C_n$ with C_n in the region $\lambda < 0$. Recall that a gallery is a chain of chambers with successive chambers sharing a common wall. The depth is also the smallest n with

$$\lambda = s_1 s_2 \dots s_{n-1} \alpha$$

for some $\alpha \in \Delta$. The elements of Δ itself, for example, have depth 1. A partial order is thus induced on the set of all roots: $\lambda \preceq \mu$ if $\mu = w\lambda$ where $\delta(\mu) = \ell(w) + \delta(\lambda)$. This order gives rise to the **root graph**, whose nodes are positive roots, with edges $\lambda \rightarrow s\lambda$ whenever $\lambda \prec s\lambda$.

1.5. Lemma. *If $\lambda > 0$ and $s_\alpha \lambda$ dominates α then $\lambda \prec s_\alpha \lambda$.*

PROOF. If (C_i) is a gallery from C to the region where $s_\alpha \lambda < 0$, there will be some pair C_i, C_{i+1} sharing a wall on the hyperplane $\alpha = 0$. The reflected gallery will be shorter. Hence $\delta(\lambda) < \delta(s_\alpha \lambda)$. $\square$

The minimal roots lie at the bottom of the root graph, in the sense that:

1.6. Proposition. (B & H 2.2(iii)) *If $\lambda \preceq \mu$ and μ is minimal, so is λ .*

PROOF. This reduces to the case $\mu = s_\alpha \lambda$. If μ is minimal but $\mu \neq \alpha$, then either $s_\alpha \mu = \lambda$ is minimal, or it dominates α . But by the preceding Lemma, in the second case $\delta(\mu) < \delta(s_\alpha \mu)$, a contradiction. $\square$

1.7. Lemma. (B & H 2.3) *If $\lambda > 0$ and $\lambda \bullet \alpha > 0$ then $s_\alpha \lambda \prec \lambda$.*

PROOF. Suppose $w^{-1}\lambda < 0$ with $\ell(w) = \delta(\lambda)$. If $w^{-1}\alpha < 0$ then we can write

$$\begin{aligned} w &= s_\alpha s_2 \dots s_n \\ \lambda &= s_\alpha s_2 \dots s_{n-1} \alpha_n \\ s_\alpha \lambda &= s_2 \dots s_{n-1} \alpha_n \\ \delta(s_\alpha \lambda) &= n-1 < \delta(\lambda) = n. \end{aligned}$$

So we may assume that $w^{-1}\lambda < 0$ but $w^{-1}\alpha > 0$. Let w have the reduced expression $w = s_1 \dots s_n$, and let $u = ws_n = s_1 \dots s_{n-1}$. Thus $w\alpha_n < 0$. I claim that $u^{-1}s_\alpha \lambda < 0$, which implies that $\delta(s_\alpha \lambda) = n-1$.

We can calculate that

$$\begin{aligned} w^{-1}s_\alpha \lambda &= w^{-1}\lambda - 2(\lambda \bullet \alpha)w^{-1}\alpha \\ u^{-1}s_\alpha \lambda &= u^{-1}\lambda - 2(\lambda \bullet \alpha)s_n w^{-1}\alpha. \end{aligned}$$

From the first equation, we see that $w^{-1}\lambda$ will be negative under the assumption that $\lambda \bullet \alpha > 0$. Since the depth of λ is n , $\mu = u^{-1}\lambda$ will be positive. But then $\mu > 0$, $s_n \mu < 0$ implies that $\mu = \alpha_n$. So we have

$$u^{-1}s_\alpha \lambda = \alpha_n - 2(\lambda \bullet \alpha)s_n w^{-1}\alpha.$$

The root

$$\alpha_n - 2(\lambda \bullet \alpha)s_n w^{-1}\alpha$$

is either positive or negative. In the second term, the positive root $w^{-1}\alpha$ cannot be α_n , since then $w\alpha_n < 0$. Therefore $s_n w^{-1}\alpha$ is a positive root also not α_n , and the second term is a negative root not equal to a multiple of $-\alpha_n$. The sum has to be negative. $\blacksquare$

1.8. Corollary. (B & H 1.7) *Whenever λ is a positive root and s in S there are three possibilities: (a) $s\lambda \rightarrow \lambda$, (b) $s\lambda = \lambda$, or (c) $\lambda \rightarrow s\lambda$, depending on whether $\lambda \bullet \alpha_s > 0$, $\lambda \bullet \alpha_s = 0$, or $\lambda \bullet \alpha_s < 0$, respectively.*

1.9. Corollary. (B & H 1.8) *If $\lambda = \sum \lambda_\alpha \alpha \preceq \mu = \sum \mu_\alpha \alpha$ then $\lambda_\alpha \leq \mu_\alpha$.*

These results will soon be used to outline an algorithm to construct the minimal root reflection table. Processing roots will take place in order of increasing depth, applying this criterion: if $\lambda \neq \alpha$ is a minimal root, then $s_\alpha \lambda$ is no longer a minimal root if and only if $\lambda \bullet \alpha_s \leq -1$. Details follow, after I explain in the next section how to use the minimal root reflection table in multiplication.

2. Minimal roots and multiplication

Suppose that w has the normal form $w = s_1 \dots s_n$. Then the normal form of sw will be obtained from that of w by **insertion** or **deletion**:

$$NF(sw) = s_1 \dots s_i t s_{i+1} \dots s_n$$

or

$$NF(sw) = s_1 \dots s_{i-1} s_{i+1} \dots s_n .$$

How do we find where the insertion or deletion occurs, and which of the two it is? If insertion, what is inserted?

The algorithm explained in (Casselman, 2001) handles these problems in geometric terms. I define the **InverseShortLex tree** to be the directed graph whose nodes are elements of W , with a link from x to y if $NF(y) = NF(x) \bullet s$. Its root is the identity element. Each $y \neq 1$ in W is the target of exactly one link, labeled by the s in S least with $ys < y$. Paths in this tree starting at 1 match the normal forms of elements of W .




Figure 1. *The InverseShortLex tree for $\tilde{A}_2$.*

This tree gives rise to a geometrical figure in which we put a link from xC to yC if there is an edge in the tree from x to y . The multiplication problem can now be formulated: *given an InverseShortLex path from C to wC , how can we find the InverseShortLex path from C to swC ?* As explained in (Casselman, 2001), the InverseShortLex figure is very close to being symmetric with reflection to the reflection s ; only links near the hyperplane $\alpha_s = 0$ are not preserved upon reflection. The chambers principally affected are what I call **exchange sites** of the InverseShortLex tree— wC is an exchange site if the chamber wC has a face on the hyperplane $\alpha_s = 0$, labeled by t in S which is less than the labelling of the InverseShortLex link entering wC .




Figure 2. Some of the exchange nodes for reflection by $\langle 1 \rangle$ in $\tilde{A}_2$. Links modified under reflection are colored. The fundamental chamber is at the bottom.

The relationship between this notion and multiplication is simple. Suppose $NF(w) = s_1 \dots s_n$ and for each i let w_i be the partial product $s_1 \dots s_i$. Suppose that $w_k C$ is the last exchange site in the gallery $C = C_0, \dots, C_n = w_n C$, and suppose that t labels the wall on $\alpha = 0$. Then either the gallery crosses $\alpha = 0$ there or it does not. In the first case, $t = s_{k+1}$ and

$$NF(sw) = s_1 \dots s_k s_{k+2} \dots s_n.$$

In the second,

$$NF(sw) = s_1 \dots s_k t s_{k+1} \dots s_n.$$

The question is now: *How do we recognize InverseShortLex exchange sites?* This is where roots and, even more pleasantly, minimal roots come in.

2.1. Lemma. Suppose $w = s_1 \dots s_n$. The chamber wC is an InverseShortLex exchange site if and only if $w^{-1}\alpha = \beta$ with $s_\beta < s_n$.

PROOF. The wall of wC labeled by β is embedded in the hyperplane $w\beta = 0$. $\square$

This suggests the following algorithm to calculate $NF(sw)$: Read the string $s_1 \dots s_n$ from left to right. As we read, keep track of an element t in S , an index k , and a root λ . The index k is that of the last exchange, the element t is the label of the last exchange, $\lambda = w^{-1}\lambda$. To start with $k = -1$, $t = s$, $\lambda = \alpha_s$. As we read s_i , λ changes to $s_i\lambda$. If $\lambda = \beta$ with $s_\beta < s_i$, t changes to s_β and k to i . When we are through scanning, we have the data determining the last exchange.

So far, minimal roots have not come into play. But in fact, in calculating the reflected root $s\lambda$ we only need to apply the minimal root reflection table. If λ is minimal and $s\lambda = \ominus$, then the chambers $w_{i-1}C_{i-1}$ and $C_i = w_{i-1}s_iC_i$ lie on opposite sides of the hyperplane $\alpha = 0$, so we know we are looking at a deletion. If $s\lambda$ is minimal and $s\lambda = \oplus$ then we know the gallery has crossed a hyperplane $\lambda = 0$ which separates wC from $\alpha = 0$. The exchange

we last recorded will be the last we ever record, because the gallery cannot recross that hyperplane to reach $\alpha = 0$.

We now know how to use the minimal root reflection table to multiply, and it remains to tell how to construct it.

3. A rough guide to constructing the minimal roots

A straightforward procedure to list all minimal roots, along with a description of how the elements of S act on them, is not difficult to sketch.

The data determining a minimal root λ are an index, a coordinate vector, and a list of reflections $s\lambda$. There are at least two reasonable choices of coordinates. The coordinates I use in this note are those in the expression of a root as a linear combination of the roots in Δ :

$$\lambda = \sum_{\alpha \in \Delta} \lambda_{\alpha} \alpha$$

The other reasonable choice is the array $(\lambda \bullet \alpha)$, which du Cloux has discovered to be a choice somewhat more efficient if also more difficult to work with in programming. Indices are assigned in the order in which the minimal roots are first calculated; the roots in Δ are assigned the first $|S|$ integers. Furthermore, in the preliminary version of the algorithm we shall maintain a look-up table of some kind, enabling us to find a root's data given its coordinates.

A minimal root is to be dealt with in two stages: (1) **defining** and (2) **finishing**. In the definition, a root is calculated, assigned an index, entered in the look-up table, and put on a waiting list. Items are taken off this list in the order in which they are put on—the list is, in programmers' terminology, a FIFO list (**first in, first out**) or **queue**.

A root λ is finished when it is taken off the queue and all of the reflections $s\lambda$ not already known are found. At this moment, all roots with depth less than that of λ have been finished, so that in particular we have recorded all reflections $\mu = s\lambda \prec \lambda$. We must calculate the $s\lambda$ not yet determined; the problem is to figure out whether $s\lambda = \lambda$, $s\lambda$ is a minimal root, or $s\lambda = \oplus$. Of course we can tell by inspection whether $s\lambda = \lambda$. Otherwise we know that $\lambda \prec \mu = s\lambda$. There are three possibilities: (1) We have already defined μ , in which case it will be registered in our look-up table. We just add to its record that $s\lambda = \mu$ and $s\mu = \lambda$. (2) The root μ is not minimal. We can tell this when we calculate μ , since in this case $\lambda \bullet \alpha_s \leq -1$. We set $s\lambda = \oplus$. (3) Otherwise $-1 < \lambda \bullet \alpha_s < 0$, and the root μ is a new minimal root not yet defined. We do so by assigning it an index, putting it in the look-up table, recording the reflection $s\lambda = \mu$, and putting μ in the process queue. Then we go on to the next root in the queue, as long as there is something there.

There are a number of problems that occur in trying to implement this in practice. The first is that maintaining the look-up table is somewhat cumbersome. As du Cloux first observed, one of the results from Brink's thesis allows us to dispense entirely it. The following is an endsatz of the main Theorem of (Brink, 1995).

3.1. Lemma. *Suppose $s \neq t$ in S . Assume the group $W_{s,t}$ they generate to be finite. Suppose λ to be any root whose $W_{s,t}$ -orbit doesn't intersect Δ . Then $W_{s,t}$ acts freely on this orbit.*

How can this be applied in our algorithm to find all minimal roots? Suppose we are in the process of defining a minimal root μ , which we may assume not to be dihedral. We calculate it as $\mu = s\lambda$ with $\lambda \prec \mu$. If t is an element of S with $t\lambda \prec \lambda$, then $W_{s,t}$ will necessarily be finite. In the orbit of μ under $W_{s,t}$ there will be a unique element ν of least depth, equal to

$w_{s,t}\mu$ where $w_{s,t}$ is the unique longest element of the dihedral group $W_{s,t}$. We can calculate it by finding alternately $t\lambda$, $st\lambda$, etc. until the chain starts to ascend in depth. And the difference in depth between μ and ν will be exactly $m_{s,t}$ precisely when $t\mu \prec \mu$. The element $t\mu$ will then be $(tw_{s,t})\nu$, which can be calculated from the reflection tables of elements of depth less than λ . In other words, as soon as μ is defined we can calculate all the **descents** $t\mu \prec \mu$ —without doing any new calculations!

In the algorithm above we can therefore eliminate reference to the look-up table. The procedure sketched above requires, however, that we know whether the reflection of a root has less depth or not, so we add to the data of a root λ its **descent set**, the set of s with $s\lambda \prec \lambda$. When we define a root, before we put it in the queue, we calculate all of its descents.

Because the result above is applicable only to roots not in the orbit of $W_{s,t}$, all the dihedral roots have to be handled somewhat specially—they should all be defined immediately after the roots in Δ , and those of depth greater than 2 have to be fed into the queue at the right moment.

There is another simplification. The following was observed by Brink and Howlett, and used as an important part of their proof that the set of minimal roots is finite:

3.2. Proposition. *If $s\lambda = \oplus$ and $\lambda \prec \mu$ then $s\mu = \oplus$ as well.*

PROOF. For α, β in Δ , λ any root

$$s_\alpha \lambda \bullet \beta = (\lambda - 2(\lambda \bullet \alpha)\alpha) \bullet \beta = (\lambda \bullet \beta) - 2(\lambda \bullet \alpha)(\alpha \bullet \beta)$$

which means that for $\beta \neq \alpha$ the dot product $\lambda \bullet \beta$ decreases under reflection by α , as long as $\lambda \prec s_\alpha \lambda$. Therefore once we have calculated that $s\lambda = \oplus$ this remain true for all μ with $\lambda \prec \mu$. $\blacksquare$

I call an s with $s\lambda = \oplus$ a **lock** at s . This Proposition tells us that locks are inherited as we go up the root graph and these inherited locks, like descents, can be immediately recorded upon definition.

Eliminating the look-up table is a great simplification, noting inherited locks a minor one. In simple cases—when none of the Coxeter matrix entries $m_{s,t}$ are greater than 3—there is probably no better algorithm than the one sketched above, but incorporating the simple modifications concerning descents and locks. In this case, all doubled dot products $2(\lambda \bullet \alpha)$ will be integers. When $\lambda \prec s_\alpha \lambda$ this doubled dot-product will be a negative integer, and if it is anything other than -1 the root $s\lambda$ will no longer be minimal. What could be simpler?

If there are entries $m_{s,t} > 3$ this still works, at least in principle, although there are a few annoyances that arise. All dot products in the range $(-1, 0)$ will be of the form $\cos(\pi k/m_{s,t})$ for $k < m_{s,t}/2$. So in order to tell whether $s\lambda$ dominates α_s we just have to compare the dot product $\lambda \bullet \alpha_s$ to numbers in this finite set. There is one potentially nasty problem, however. All we can assert *a priori* about the coordinates of λ is that they will be in the real cyclotomic field generated by $\cos(2\pi/M)$, where M is the lowest common denominator of the numbers $2m_{s,t}$. The amount of work involved would seem to grow disturbingly as M grows. It is Brink's thesis that tells us that this apparent difficulty is spurious. In the next section I follow her in examining more carefully the coefficients of minimal roots.

4. Root coefficients

We want to know something about what possible values can occur for the coefficients of roots and, especially, minimal roots. Roots are produced by a sequence of reflections from the basic roots in Δ . As is the case in many algorithmic processes, although we understand each single step—here, reflection—quite well, the overall development is not so clear.

We begin with a discussion of dihedral roots.

Suppose S to possess two elements s and t , corresponding to roots α and β . Let $m = m_{s,t}$. What are all the positive roots of the system?

Let $z = \zeta_{2m} = e^{\pi i/m} = e^{2\pi i/2m}$. Then $2\alpha \bullet \beta = 2\cos(\pi/m) = (z + z^{-1})$ and

$$\begin{aligned} s\lambda &= \lambda - 2(\lambda \bullet \alpha)\alpha \\ 2(p\alpha + q\beta) \bullet \alpha &= 2p - (z + z^{-1})q \\ s(p\alpha + q\beta) &= [-p + (z + z^{-1})q]\alpha + q\beta \\ t\lambda &= \lambda - 2(\lambda \bullet \beta)\beta \\ 2(p\alpha + q\beta) \bullet \beta &= 2q - (z + z^{-1})p \\ t(p\alpha + q\beta) &= p\alpha + [-q + (z + z^{-1})p]\beta. \end{aligned}$$

Suppose $m = \infty$. We get in succession (in tabular form) as transforms of α

	coefficient of α	coefficient of β
α	1	0
$t\alpha$	1	2
$st\alpha$	3	2
$\dots$	$\dots$	$\dots$
$(st)^n\alpha$	$2n + 1$	$2n$
$t(st)^n\alpha$	$2n + 1$	$2n + 2$
$\dots$		

and similarly for the transforms of β . Of these, only α and β are minimal roots.

If $m = 2$, the positive roots are α and β .

Now suppose $2 < m < \infty$. Then we get

	coefficient of α	coefficient of β
α	1	0
$t\alpha$	1	$z + z^{-1}$
$st\alpha$	$z^2 + z^{-2} + 1$	$z + z^{-1}$
$\dots$	$\dots$	$\dots$
$(st)^n\alpha$	C_{n+1}	C_n
$t(st)^n\alpha$	C_{n+1}	C_{n+2}

where the C_n are solutions of the difference equation $C_{n+2} = -C_n + (z + z^{-1})C_{n+1}$ with initial values $C_0 = 0$, $C_1 = 1$, giving

$$C_n = \frac{z^n - z^{-n}}{z - z^{-1}} = z^{n-1} + z^{n-3} + \dots + z^{-(n-3)} + z^{-(n-1)} = \frac{\sin(\pi n/m)}{\sin(\pi/m)}.$$

Note that $C_n > C_{n-1}$ as long as the real part of z^n is positive, which happens as long as $n < m/2$. There are m roots in all for a dihedral system, which agrees with this remark.

For $m = 3$ we get the table

	coefficient of α	coefficient of β
α	1	0
β	0	1
$t\alpha = s\beta$	1	1

For $m = 4$:

	coefficient of α	coefficient of β
α	1	0
β	0	1
$t\alpha = st\alpha$	1	$\sqrt{2}$
$s\beta = ts\beta$	$\sqrt{2}$	1

For $m = 5$, with $\theta = (1 + \sqrt{5})/2$:

	coefficient of α	coefficient of β
α	1	0
β	0	1
$t\alpha$	θ	1
$s\beta$	1	θ
$st\alpha = ts\beta$	θ	θ

For $m = 6$:

	coefficient of α	coefficient of β
α	1	0
β	0	1
$t\alpha = st\alpha$	1	$\sqrt{3}$
$s\beta = ts\beta$	$\sqrt{3}$	1
$st\alpha = tst\alpha$	2	$\sqrt{3}$
$ts\beta = sts\beta$	$\sqrt{3}$	2

The table for $m > 6$ looks somewhat similar to the last, except that it is longer. All coefficients beyond the first 6 rows are greater than 2.

We can use these results, following Brink's thesis, to prove:

4.1. Proposition. *The coefficients of the positive roots are polynomials in the constants*

$$\frac{\zeta_{2m}^n - \zeta_{2m}^{-n}}{\zeta_{2m} - \zeta_{2m}^{-1}} \quad (n \leq m/2)$$

with coefficients among the natural numbers $\mathbb{N}$.

PROOF. By induction on the depth of a root. If the depth is 1 there is no problem. Otherwise suppose

$$\lambda = w\alpha = s_1 \dots s_{n-1}\alpha_t$$

with α in Δ , $\delta(\lambda) = n$. Let $s = s_n$. Choose $y \neq 1$ maximal in $W_{s,t}$ with $w = xy$. Then $x\alpha_s > 0$, $x\alpha_t > 0$. We know from the discussion of dihedral roots what the coefficients of

$$y\alpha_t = p\alpha_s + q\alpha_t$$

are like, but then

$$w\alpha_t = px\alpha_s + qx\alpha_t.$$

We can apply an induction hypothesis to $x\alpha_s$ and $x\alpha_t$ since $\ell(x) < \ell(w)$. $\square$

4.2. Corollary. *The smallest positive root coefficient is 1. Any root coefficient lying between 1 and 2 equals $2 \cos(\pi/m_{s,t})$ for some s, t .*

PROOF. If

$$\frac{\sin(\pi n/m)}{\sin(\pi/m)} \quad (1 \leq n \leq m/2)$$

lies between 1 and 2 then $n = 2$, and the least possible value is $\sqrt{2}$. Any root coefficient will be a sum of terms

$$C = c_1 \dots c_k$$

where $k \geq 1$ and each c_i will be either a positive integer or one of the constants in the Proposition, with each $c_i > 1$. The smallest possible value for one of these constants is $2 \cos(\pi/4) = \sqrt{2}$, so if this product is less than 2 then $k = 1$, and is an integer only if the product itself, of course, is 1. $\square$

4.3. Corollary. *The smallest possible coefficient of a positive root, other than 1, is $\sqrt{2}$.*

5. Composition and decomposition

One of the important results of Brink's thesis is that the apparent difficulties concerning mixed cyclotomy in the determination of the minimal roots are in fact spurious. In order to see that it is true, we have to look carefully at how roots are calculated.

A root $\sum \lambda_\alpha \alpha$ may be considered as a function on the Coxeter graph: $s \mapsto \lambda_{\alpha_s}$. The **support** of a root $\lambda = \sum \lambda_\alpha \alpha$ is its support as a function—the set of s where $\lambda_{\alpha_s} \neq 0$. Brink's nice idea is to track the computation of roots by looking at their support. All roots are constructed by applying some w to a basic root. The support of a root λ is extended by applying a reflection s which is not already in its support. If s is not connected to that support in the Coxeter graph, then $s\lambda = \lambda$. This process gives a new root only when s is linked to the support of λ . Therefore:

5.1. Proposition. *The support of any root is connected.*

Suppose we apply a reflection s to a positive root λ whose support does not contain s . Let $\alpha = \alpha_s$. Only the coefficient λ_α changes, and to

$$-\lambda_\alpha + \sum_{\lambda_\beta \neq 0, \alpha \sim \beta} [-2\alpha \bullet \beta] \lambda_\beta.$$

If λ is a minimal root with $\lambda_{\alpha_s} = 0$, then $s\lambda$ will also be a minimal root if and only if

$$\sum_{\lambda_\beta \neq 0, \beta \sim \alpha} [-2\alpha \bullet \beta] \lambda_\beta \leq 2.$$

But $\lambda_\beta \geq 1$ and also $-2\alpha \bullet \beta \geq 1$. The condition for minimality cannot hold here unless there is just one $\beta = \alpha_t$ in the sum. Furthermore, in that case the above condition amounts to the condition

$$[-2\alpha \bullet \beta] \lambda_\beta \leq 2.$$

The first term is either 1 or $\geq \sqrt{2}$, as is the coefficient λ_β . All in all:

5.2. Proposition. *Suppose λ to be a minimal root, s an element of S not in the support Θ of λ . If $s\lambda$ is again minimal then there must be just a single link from s to Θ , say to t , and one of these holds:*

- (a) $m_{s,t} = 3$ and $\lambda_{\alpha_t} < 2$;
- (b) $3 < m_{s,t} < \infty$ and $\lambda_{\alpha_t} = 1$.

We can summarize this in a mnemonic diagram:

$$\begin{aligned}
 \bullet \quad x \text{---} 0 &\longmapsto x \text{---} x \quad (x < 2) \\
 \bullet \quad 1 \text{==} 0 &\longmapsto 1 \text{==} c \quad (c = 2 \cos(\pi/m))
 \end{aligned}$$

5.3. Corollary. *The support of a minimal root is a tree with containing no links of infinite degree.*

One of Brink's good ideas is to focus on the coefficients of a minimal root which are equal to the minimum possible value 1. Suppose λ to be a minimal root with $\lambda_\alpha = 1$. We know that λ has been built up from a chain of minimal root predecessors

$$\lambda_1 \prec \dots \prec \lambda_n = \lambda.$$

Some one of these, say $\mu = \lambda_k$, will be the first to have a non-zero coefficient at α , and that coefficient must be 1, since it can only grow. Unless $\mu = \lambda_1 = \alpha$, the root μ must have been constructed by extension from a root λ_{k-1} whose support did not contain s , and the earlier Proposition tells us that $\mu = s\lambda_{k-1}$, where s is linked to the support of λ_{k-1} at a single node t by a simple link in the Coxeter graph, and $\lambda_{k-1, \alpha_t} = \lambda_{k, \alpha_t} = 1$. We have $\lambda = w\mu$ where w is a product of generators other than s .

Since the support Θ of λ is a tree, the complement of α in Θ is the disjoint union of exactly as many components $\Theta_i^\bullet$ as there are edges in the support of λ linked to α , say ℓ . Reflections corresponding to nodes in different components will commute. Hence we can write $\lambda = w_1 \dots w_\ell \mu$ where each w_i is in $W_{\Theta_i^\bullet}$. All the w_i commute, so we can write the product in any order. Each of the $\lambda_i = w_i \alpha$ is itself a root with support in $\Theta_i = \Theta_i^\bullet \cup \{\alpha\}$, minimal because it will be a predecessor of λ in the root graph. We have therefore proven:

5.4. Proposition. *Let λ be a minimal root with $\lambda_\alpha = 1$. Let Θ be the support of λ , the $\Theta_i^\bullet$ the connected components of $\Theta \setminus \{\alpha\}$, $\Theta_i = \Theta_i^\bullet \cup \{\alpha\}$. The restriction λ_i of λ to Θ_i is also a minimal root.*

This is called the **decomposition** of λ . Conversely:

5.5. Proposition. *Let Θ be a subset of S , α an element of Θ , the $\Theta_i^\bullet$ the components of $\Theta \setminus \{\alpha\}$, $\Theta_i = \Theta_i^\bullet \cup \{\alpha\}$. If for each i the minimal root λ_i has support on Θ_i with coefficient 1 at α , then the vector λ with support Θ whose restriction to each Θ_i agrees with λ_i is a minimal root.*

This is called **composition**.

PROOF. More difficult than the previous result. It follows from this result of Brink's:

5.6. Lemma. *If λ is any positive root with coefficient 1 at α , then $\lambda = w\alpha$, where w is a product of s in S not equal to s_α .*

PROOF. Induction on the depth of λ . Trivial if $\lambda = \alpha$. It remains to be shown that if $\delta(\lambda) > 1$ and $\lambda_\alpha = 1$ then we can find s such that $s\lambda \prec \lambda$ and $s\lambda$ also has coefficient 1 at α . There exists at any rate some s with $s\lambda \prec \lambda$. If this does not have coefficient 1 at α , then—as we have seen above— $s\alpha$ has coefficient 0 at α ; α has exactly one link to the support of $s\lambda$, say to β ; and the coefficient of $s\lambda$ at β is equal to 1. By induction, $s\lambda = w\beta$ where w is a product of t equal neither to s nor s_β . But then

$$\lambda = sw\beta = ws\alpha \quad \square$$

These results have a natural generalization. The **unit support** of λ is the set of α with $\lambda_\alpha = 1$. Suppose λ to be a minimal root, Θ its support, $T \subset \Theta$ its unit support. We know that Θ is a connected tree. The complement $\Theta \setminus T$ can be expressed as the disjoint union of its connected components $\Theta_i^\bullet$. For each of these, let Θ_i be its union with elements of T attached to it. The set Θ_i will then be connected, and no point of T will be in its interior. The set T will contain in addition some pairs of nodes of the Coxeter graph connected by links in it; let these be counted as additional sets Θ_i (but with empty interiors). Finally, there is one exceptional case where Θ has just one element, which is Θ_1 .

5.7. Lemma. *The set Θ is the union of the Θ_i , which overlap only in points of T .*

PROOF. We may suppose Θ to have more than one element in it. Suppose given t in Θ . If it is not in T , it belongs to a unique $\Theta_i^\bullet$. Otherwise, it will be in T . If it is connected in the Coxeter diagram to one of the $\Theta_i^\bullet$, it will be in Θ_i . Finally, we have the situation where the only elements of Θ it is linked to are in T . Each of the pairs forming such a link will be one of the Θ_i . $\square$

If T is any subset of S , a subset Θ is called **T-connected** if none of its interior points belong to T . Each of the Θ_i defined above are T -connected, and are called its **T-components**. Following Brink, I call a minimal root **indecomposable** if its unit support is contained in the boundary of its support.

Lemma. (Brink) (Decomposition) *The restriction of a minimal root to any of the T -components of its support is again a minimal root.*

PROOF. This follows by induction from the earlier decomposition. $\square$

5.8. Proposition. (Brink)(Composition) *Let Θ be any connected, simply connected subset of S , T an arbitrary subset of Θ , Θ_i its T -components. If λ_i is for each i a minimal root with support Θ_i and unit support $T \cap \Theta_i$, then the vector λ with support Θ whose restriction to Θ_i is λ_i will be a minimal root.*

This reduces the calculation of minimal roots to the calculation of indecomposable ones, together with a bit of accounting. Finding the reflection table turns out not to be much more difficult.

How are indecomposable roots found? They are obtained by reflection from predecessors in the root graph. What can we say about this process? If λ is an indecomposable root and $\mu = s_\alpha \lambda \prec \lambda$, what can we say about μ ? Let Θ be the support of λ . There are several possibilities. (1) Suppose that α is a boundary point of Θ , and that $\lambda_\alpha = 1$. Then λ is obtained from μ by **extension** from a unit node of μ . This must be a boundary point of the support of μ . This can happen only if μ has singleton support. (2) Suppose that α is in the boundary of Θ but that $\lambda_\alpha > 1$. Then μ will also be indecomposable. (3) Suppose α lies in the interior of Θ and $\mu_\alpha > 1$. Then μ is also indecomposable. (4) Finally, suppose α lies in the interior of Θ but $\mu_\alpha = 1$. The support of μ is also Θ , so μ cannot be indecomposable. It will be the composite of several μ_i adjoined to each other at α . Thus:

5.9. Proposition. *An indecomposable root is constructed through a sequence made up of three types of processes: (1) **extension**, where $\mu \prec \lambda = s_\mu \mu$, μ is also indecomposable, and the support of λ is one node larger than the support of μ ; (2) **fusion**, where μ is the composite of indecomposable roots μ_i all with α in the unit support; (3) **promotion**, where μ is indecomposable and has the same support as λ .*

We have already investigated extension. In the next section we look at fusion.

6. Reflecting at junctions

Suppose Θ to be a connected and simply connected subset of S , $\alpha \in \Theta$, the Θ_i the connected components of $\Theta \setminus \{s\}$ joined with $\{s\}$. Suppose that λ is a minimal root with support Θ and $\lambda_{\alpha_s} = 1$. Let λ_i be the restriction of λ to Θ_i , and assume the λ_i are indecomposable. *Under what circumstances is $s\lambda$ a minimal root?*

Each of the Θ_i contains a unique node t_i linked to s . Let $\beta_i = \alpha_{t_i}$.

The inner product $\lambda \bullet \alpha$ is

$$1 - \sum \lambda_{\beta_i} |\alpha \bullet \beta_i|.$$

Recall that $\lambda_{\beta_i} \geq 1$, $|\alpha \bullet \beta_i| \geq 1/2$. When the junction has more than 3 roots joining at it, this dot-product is ≤ -1 , and $s\lambda$ will not be minimal.

6.1. Proposition. *If $\Theta \setminus \{s\}$ has more than 3 components, $s\lambda = \oplus$.*

We may now assume there to be 2 or 3 links at α .

6.2. Lemma. *Suppose $c = 2 \cos(\pi/m_{s,t})$. The links $c \text{---} 1$ and $1 \text{==} 1$ cannot occur in any root.*

This introduces notation I hope to be self-explanatory. What I mean by this is that if λ is a root then there do not exist neighbouring nodes s and t in the Coxeter graph with (1) $m_{s,t} = 3$, $\lambda_{\alpha_s} = 1$, $1 < \lambda_{\alpha_t} < 2$; or (2) $m_{s,t} > 3$, $\lambda_{\alpha_s} = 1$, $\lambda_{\alpha_t} = 1$.

PROOF. By induction. The only way to get $c \text{---} 1$ is from $c \text{---} 1 \text{---} 1$. $\square$

We now consider the possibilities for fusion.

- *Two links.*

The three *a priori* possibilities are

$$\begin{array}{c} \lambda_x \text{---} 1 \text{---} \lambda_y \\ \lambda_x \text{---} 1 \text{==} \lambda_y \\ \lambda_x \text{==} 1 \text{==} \lambda_y \end{array}$$

6.3. Proposition. *In the third case $s\lambda = \oplus$.*

PROOF. The inner product is $1 - c_x \lambda_x / 2 - c_y \lambda_y$. We know that each c_* is at least $\sqrt{2}$, and by the previous Lemma that $\lambda_* \geq \sqrt{2}$ as well. Thus the dot product is -1 or less. $\square$

In the first case, the condition is $1 - \lambda_x / 2 - \lambda_y / 2 > -2$, or $\lambda_x + \lambda_y < 4$. The case $\lambda_* = c$ is excluded, so either $\lambda_* = 1$ or $\lambda_* \geq 2$. It is not possible for both λ_* to be ≥ 2 , so one at least must be 1, say $\lambda_y = 1$.

We can do better: the inner product must be $-\cos(\pi k/m)$ for $k \leq m/2$. This gives us:

$m = 3$. The inner product is 0, with

$$\bullet \quad 1 \text{---} 1 \text{---} 1 \quad \mapsto \quad 1 \text{---} 1 \text{---} 1$$

or $-1/2$ with

$$\bullet \quad 2 \text{---} 1 \text{---} 1 \quad \mapsto \quad 2 \text{---} 2 \text{---} 1.$$

$m = 4$. We must have $1/2 - \lambda_x / 2 = -\sqrt{2}/2$ or $\lambda = 1 + \sqrt{2}/2$. Impossible.

$m = 5$. $1 - \lambda_x = -c$ or $1 - c$, $\lambda_x = 1 + c$ or $\lambda = c$. The second is impossible, alas. So the only possibility is

$$\bullet \quad (1+c) \longrightarrow 1 \longrightarrow 1 \quad \mapsto \quad (1+c) \longrightarrow (1+c) \longrightarrow 1.$$

$m \geq 6$. Nothing possible.

Now the remaining case

$$\lambda_x \longrightarrow 1 \rightleftharpoons \lambda_y,$$

Here

$$1 - \lambda_x/2 - (c/2)\lambda_y > -1, \quad \lambda_x + c\lambda_y < 4.$$

$m = 4$. Only possibility is

$$\bullet \quad 1 \longrightarrow 1 \rightleftharpoons \sqrt{2} \quad \mapsto \quad 1 \longrightarrow 2 \rightleftharpoons \sqrt{2}.$$

$m = 5$. Only possibility is

$$\bullet \quad 1 \longrightarrow 1 \rightleftharpoons c \quad \mapsto \quad 1 \longrightarrow (1+c) \rightleftharpoons c.$$

Note that in all these cases we have a join of two primitive roots, one of which is just $1 \longrightarrow 1$. So it is trivial to reconstruct the reflection even when this is all embedded in a generic root.

• *Three links.*

Assume the reflection is a primitive root. The conditions are $\lambda_x + \lambda_y + \lambda_z < 4$ if all links in are simple, or $\lambda_x + \lambda_y + c_m \lambda_z < 4$ if one has degree $m > 3$. More of degree $m > 3$ cannot occur by simple calculation. If $m > 3$ then $c \geq \sqrt{2}$ so the second sum is at least 4, and cannot occur.

As for the first, we cannot have any $\lambda_* \geq 2$, so each can only be 1 or c . But the link $c \longrightarrow 1$ cannot occur.

6.4. Proposition. *The only triple junction producing a minimal root be reflection at the junction is the case that all coefficients are 1. It reflects to a central node with value 2.*




Figure 3. *The only triple junction giving rise to fusion.*

Looking back over all ways indecomposable minimal roots are constructed, we deduce:

6.5. Proposition. *No indecomposable minimal root has more than one link of degree $m > 3$ in its support.*

This means that mixed cyclotomy does not occur. Brink's thesis at this point goes on to build a rather explicit description of all possible indecomposable roots whose support is a given Coxeter graph. It is probably a poor idea to try to use these lists in a computer program, since it is fairly simple to use basic properties of indecomposable roots to build these lists automatically. There is one consequence of what we have seen so far that is simple to formulate. It turns out that any root coefficient may be described in no more than three integers: each one is either of the form $a + bc_m$ with $c_m = 2 \cos(\pi/m)$ or larger than 2. With a bit of fiddling to handle the last case carefully, the three integers (a, b, m) are sufficient. This is a drastic version of the principle of 'no mixed cyclotomy'. Implementing this idea in an explicit program is straightforward enough, but the program to be sketched in the next section, which uses composition and decomposition in a more sophisticated fashion, seems to be more efficient.

There is one final point to take into account. When an indecomposable root λ has degree $m > 6$, it will have a unique link in its support of that degree. The discussion of extension above tells us that this link is in the support of the dihedral root first encountered in constructing λ . Suppose this dihedral root has support at s and t , and that it started with s . The discussion of dihedral roots tells us that the coefficient of α_s in λ will be either 1 or greater than 2. In the first case, s has to be on the boundary of the support of λ because of indecomposability. In the second, we know that it is on the boundary because no node with coefficient ≥ 2 can be extended. In all cases, therefore, s is on the boundary of the support of λ . I leave it as an exercise to show further that *in the second case the coefficient of α_t is equal to $c_m = 2 \cos(\pi/m_{s,t})$, and in both cases all other coefficients are of the form nc_m where n is a positive integer.* (This, too, is an observation of Brink's thesis.)

7. The program

I am going to sketch here a program that lists all minimal roots and constructs the reflection table. Relying strongly on Brink's thesis, it works with two kinds of root, **indecomposable** and **composite**. The associated data structures are quite different. Both have indices, descent sets, and a reflection list. But an indecomposable root has in addition an actual array of coefficients. These coefficients will all lie in one of the real cyclotomic rings generated by $\zeta_{2m} + \zeta_{2m}^{-1}$, where $m = m_{s,t}$ for some s, t . I call these rings **elementary**. An indecomposable root is also assigned its degree m , and when $m > 6$ its two special nodes, the ones spanning the unique link of degree m , are specified as well. A composite root has, instead, a list of its indecomposable components.

New roots are found by applying a reflection to a root already defined. There are several basic ways in which to do this: (1) **extension** of an indecomposable root to produce a new indecomposable root with a larger support; (2) **promotion** of an indecomposable root to produce a new indecomposable root with the same support; (3) **fusion** at a junction of several indecomposable roots to produce a new indecomposable root; (4) **composition** of an arbitrary root with a dihedral root attached at one end to its unit support; (5) **replacement** of one or more of the components of a composite root by a single indecomposable root.

It is only the first three that require actual arithmetic, and always in one of the elementary real cyclotomic rings. Composition is an extremely simple operation, while replacement amounts to reducing a composite reflection to one of the first three. That this can be done easily seems almost an accident depending on the limited number of ways fusion can take

place. I'll not give details of this below, but I'll sketch an example here. Suppose we want to calculate $s_3\lambda$ where

$$\lambda: 1 \text{---} 1 \text{---} 1 \text{---} \sqrt{2} \text{---} \sqrt{2}$$

and $s = s_3$ is the third node from the left. The root λ is the composite of three indecomposable roots

$$1 \text{---} 1, \quad 1 \text{---} 1, \quad 1 \text{---} \sqrt{2} \text{---} \sqrt{2}$$

There are two components in the **star** of s , the components of λ containing it, and they do not exhaust λ . Instead, they make up the smaller root

$$\mu: 1 \text{---} 1 \text{---} \sqrt{2} \text{---} \sqrt{2}$$

whose reflections we shall have already calculated when λ is taken off the queue. It is easy enough to locate μ , given the reflection tables already made, since it is equal to s_2 applied to the indecomposable root

$$\nu: 1 \text{---} \sqrt{2} \text{---} \sqrt{2}$$

We look up $s_2\nu$ to get μ , then look up $s\mu$, which is the indecomposable root

$$1 \text{---} 2 \text{---} \sqrt{2} \text{---} \sqrt{2}.$$

Finally, we replace the two components of μ by the single component $s\mu$ in the list of components of λ . All this can be carried out, with minor modifications, for each of the cases of fusion. In the pseudo-program below the details will be grossly telescoped.

Here is the program:

First of all we define the basic roots, those in Δ .

Then we define specially the dihedral roots, those whose support consists of exactly two nodes of the Coxeter graphs. When we are through with these we have finished the basic roots. We put the dihedral roots of depth 2 in the queue, and after we have defined all of those we put those of depth 3 in the queue. In this way, we preserve the essential property of the queue that roots of depth d are put on before those of depth $d + 1$.

As for dihedral roots of depth 4 or more, we can finish them as soon as they are constructed. Both of their coefficients are 2 or more, which means they cannot be extended, so the only reflections higher in the root graph are to dihedral roots with the same support. In other words, we never put them in the queue. Incidentally, these occur only for links with $m \geq 6$. One consequence is that in calculating reflections of roots taken off the queue, we have only to do arithmetic with algebraic integers of the form $a + bc_m$, where m is the degree of the root, $c_m = 2\cos(\pi/m)$, a and b are integers.

At this point, all roots of depth 1 have been finished, all dihedral roots have been defined, and some have been finished. The queue contains only dihedral roots. While the queue is not empty, we remove roots from it to be finished.

Let λ be a root removed from the queue. All roots of less depth have been finished already, and in particular all descents $s\lambda \prec \lambda$ have been assigned. We run through all the s in S , and for each one of them where $s\lambda$ has not been assigned, we have to calculate $s\lambda$. Exactly what this involves depends on a number of things—the basic idea is to use composition and decomposition to avoid repeating actual arithmetic.

So now suppose we are looking at s and λ with $s\lambda$ not known. Let $\alpha = \alpha_s$. There are many cases to deal with.

- The node s lies in the support of λ .
 - It belongs to a unique component λ_i .
 - That component is λ itself, which is therefore indecomposable. We must calculate the dot product $\lambda \bullet \alpha$ to see if $s\lambda$ is minimal or not, and if it is define it.
 - The component λ_i is not all of λ , which is a composite root. We have calculated $s\lambda_i$ already, and $s\lambda$ is obtained by replacing λ_i with $s\lambda_i$.
 - It belongs to several components. It is therefore a junction, and we are looking at a possible fusion. Let μ be the star of s , the composite of those components whose supports contain s .
 - The root μ is the same as λ . We have to calculate $s\lambda$ explicitly. It may be λ , $\oplus$, or a new indecomposable root.
 - The root μ makes up only a part of λ . We have calculated $s\mu$, and obtain $s\lambda$ by replacing μ with $s\mu$. Exactly what we do to find μ and $s\mu$ depends on which case of fusion we are dealing with, as I mentioned in the example I looked at earlier. This can be decided by local information around s .
- The node s does not lie in the support of λ .
 - It is linked to more than one node in the support, or by a link of infinite degree. Here $s\lambda = \oplus$.
 - It is linked by a single link of finite degree.
 - The link is to a unit node of λ . Then $s\lambda$ is a new composite.
 - The link is not to a unit node.
 - The root λ is indecomposable. We have to calculate explicitly.
 - The root λ is composite. If μ is the component it is linked to, we have already calculated $s\mu$. We replace μ by $s\mu$ in λ .

References

1. Brigitte Brink, ‘The set of dominance-minimal roots’, available as Report 94–43 from the School of mathematics and Statistics at the University of Sydney:
<http://www.maths.usyd.edu.au:8000/res/Algebra/Bri/dom-min-roots.html>
2. Brigitte Brink, ‘The set of dominance-minimal roots’, *Journal of Algebra* **206** (1998), 371–412. This publication, although it has the same title as the previous item, is very different from it. For purposes of computation it is decidedly less interesting, since it doesn’t introduce the indecomposable minimal roots, and emphasizes hand-lists instead.
3. Brigitte Brink, ‘On centralizers of reflections in Coxeter groups’, published in *Bulletin of the London Mathematical society* **28** (1996), 465–470. Also available as a preprint from
<http://www.maths.usyd.edu.au:8000/res/Algebra/Bri/>
4. Brigitte Brink and Robert Howlett, ‘A finiteness property and an automatic structure for Coxeter groups’, *Math. Ann.* **296** (1993), 179–190.
5. Bill Casselman, ‘Automata to perform basic calculations in Coxeter groups’, in *Representations of groups*, CMS Conference Proceedings 16, A.M.S., 1994.
6. Bill Casselman, ‘Computation in Coxeter groups—multiplication’, preprint, 2001. To appear in the *Electronic Journal of Combinatorics*.
7. Bill Casselman, ‘Java source code for finding minimal roots’, at
<http://www.math.ubc.ca/people/faculty/cass/java/coxeter/roots/>

-
- 8.** Fokko Du Cloux, ‘Un algorithme de forme normale pour les groupes de Coxeter’, preprint, Centre de Mathématiques à l’École Polytechnique, 1990.
 - 9.** J. Tits, ‘Le problème des mots dans les groupes de Coxeter’, *Symposia Math.* **1** (1968), 175–185.
 - 10.** E. B. Vinberg, ‘Discrete linear groups generated by reflections’, *Math. USSR Izvestia* **5** (1971), 1083–1119.